

Vacuum energy and relativistic invariance

E. Kh. Akhmedov^(a,b)

^(a)CFIF, Departamento de Fisica, Instituto Superior Técnico, P-1049-001 Lisboa, Portugal

^(b)National Research Centre Kurchatov Institute, Moscow 123182, Russia

(v1: April 4, 2002; v2: June 4, 2002)

It is argued that the zero-point energies of free quantum fields diverge at most quadratically and not quartically, as is generally believed. This is a consequence of the relativistic invariance which requires that the energy density of the vacuum ρ and its pressure p satisfy $\rho = -p$. The usually obtained quartic divergence is an artifact of the use of a noninvariant regularization which violates this relation. One consequence of our results is that the zero-point energies of free massless fields vanish. Implications for the cosmological constant problem are briefly discussed.

PACS: 11.10-z, 98.80.-k

FISIST/07-2002/CFIF

hep-th/0204048

Vacuum energy is an additive part of the energy of any physical system. It plays no role in non-gravitational physics where only energy differences are observable; however, it is of profound importance for gravity. The reason for this is that the energy itself (and not energy differences) is the source of gravity. The vacuum energy enters into the Einstein equation as a cosmological constant term. It has long been known from cosmological observations that the energy density of the vacuum ρ must be smaller than the critical density of the universe, $\rho_c \sim 10^{29} \text{ g/cm}^3 \simeq 4.3 \times 10^{-47} \text{ GeV}^4$; recent observations [1] indicate that $\rho \neq 0$ and is actually about $0.7\rho_c$. At the same time, it is frequently stated that the natural value for the vacuum energy density is $\sim M_{\text{Pl}}^4$ which is more than 120 orders of magnitude larger. This huge discrepancy between the observed and “natural” values of the vacuum energy constitutes the so called cosmological constant problem [2].

In the present Letter we argue that the above estimate of the natural value of the vacuum energy is actually incorrect and results from the use of a regularization of the zero-point energies of quantum fields that does not respect the relativistic invariance of the problem. When invariant regularizations are employed, the usual quartic divergences of the zero-point energies, which lead to the above predictions, do not arise, and these energies diverge at most quadratically. In particular, in flat spacetimes the zero-point energies of free massless fields vanish. While our observations do not solve the cosmological constant problem, we believe they may shed some new light on it.

In general, the vacuum energy gets contributions from both classical and quantum effects. Classical contributions may arise from non-vanishing effective potentials of scalar fields or scalar combinations of higher-spin fields; in particular, cosmological phase transitions may lead to “latent heat” contributions to ρ . Quantum contributions are those of the zero-point energies of all quantum fields. It is these quantum contributions to the vacuum energy that we shall be concerned with here.

Consider the zero-point energy of a free real scalar field ϕ of mass m (the generalization to the cases of a complex

scalar field or higher-spin fields is straightforward). Inserting into the energy-momentum tensor $T_{\mu\nu}$ of the field ϕ the plane-wave decomposition of this field and taking the vacuum expectation value of the component T_{00} , one arrives at the following well-known result for the contribution of the field ϕ to the energy density of the vacuum:

$$\rho = \frac{1}{2} \int \frac{d^3k}{(2\pi)^3} \sqrt{\mathbf{k}^2 + m^2}. \quad (1)$$

This is just the sum over all modes of the zero-point energies $\omega_k/2 = \sqrt{\mathbf{k}^2 + m^2}/2$. The integral in (1) is clearly divergent; introducing a 3-dimensional ultraviolet momentum cutoff Λ one finds that the integral diverges as Λ^4 . This is the usual quartic divergence of the zero-point energies discussed in many quantum field theory textbooks. Taking Λ to be the highest known energy scale, the Planck scale, one arrives at the estimate $\rho \sim M_{\text{Pl}}^4$ mentioned above.

To see what the flaw of this argument is, let us consider the contribution of the zero-point fluctuations of the field ϕ to the vacuum pressure p . First, let us recall that, quite generally, on the grounds of relativistic invariance, the energy-momentum tensor of the vacuum must be of the form

$$T_{\mu\nu}^{vac} = \text{const} \cdot g_{\mu\nu}, \quad (2)$$

where $g_{\mu\nu}$ is the spacetime metric. The vacuum pressure is given by the components T_{ii}^{vac} ($i = 1, 2, 3$) of this tensor. From eq. (2) it then immediately follows that the vacuum energy density ρ and pressure p must satisfy [3]

$$\rho = -p. \quad (3)$$

Next, we calculate the contribution of the field ϕ to the vacuum pressure. Following the same steps that led to (1) one finds

$$p = \frac{1}{6} \int \frac{d^3k}{(2\pi)^3} \frac{\mathbf{k}^2}{\sqrt{\mathbf{k}^2 + m^2}}, \quad (4)$$

where the isotropy of the vacuum has been used. It should be noted that the derivations of eqs. (1) and (4)

were actually carried out in Minkowski spacetime; we shall comment on the general case of curved spacetimes later on.

We now want to see if the relativistic invariance condition (3) is satisfied by expressions (1) and (4). Since the corresponding integrals diverge, they can only be compared after a regularization procedure has been applied. Any regularization includes, explicitly or implicitly, some subtraction; our point here is that the subtraction must be performed in such a way that the relativistic invariance condition (3) be satisfied.

We first check the standard regularization of the integrals in (1) and (4) by a 3-dimensional momentum cutoff Λ . One can write $\rho = (1/4\pi^2)I_1$, $p = (1/12\pi^2)I_2$, where

$$I_1 = \int_0^\Lambda dx x^2 \sqrt{x^2 + m^2}, \quad I_2 = \int_0^\Lambda dx \frac{x^4}{\sqrt{x^2 + m^2}}. \quad (5)$$

The relativistic invariance condition (3) implies

$$I_2 = -3I_1. \quad (6)$$

Since both I_1 and I_2 given in eq. (5) are positive definite, this condition is clearly violated. To see what kind of terms violate it, let us calculate the integral I_2 by parts. This gives

$$I_2 = \Lambda^3 \sqrt{\Lambda^2 + m^2} - 3I_1. \quad (7)$$

One sees that eq. (6) is violated by the first (off-integral) term on the r.h.s. of (7), which diverges as Λ^4 in the limit $\Lambda \rightarrow \infty$. Direct calculation of the integrals I_1 and I_2 yields

$$3I_1 = \frac{3}{4} \left[\Lambda(\Lambda^2 + m^2)^{3/2} - \frac{1}{2} \Lambda m^2 \sqrt{\Lambda^2 + m^2} - \frac{1}{2} m^4 \ln \left(\frac{\Lambda + \sqrt{\Lambda^2 + m^2}}{m} \right) \right], \quad (8)$$

$$I_2 = \frac{3}{4} \left[\frac{1}{3} \Lambda^3 \sqrt{\Lambda^2 + m^2} - \frac{1}{2} \Lambda m^2 \sqrt{\Lambda^2 + m^2} + \frac{1}{2} m^4 \ln \left(\frac{\Lambda + \sqrt{\Lambda^2 + m^2}}{m} \right) \right]. \quad (9)$$

We observe that, while the terms $\sim m^4 \ln(\Lambda)$ in these expressions satisfy the relativistic invariance condition (6), the terms of order Λ^4 and $\Lambda^2 m^2$ do not. Similar results are obtained if, instead of using a sharp cutoff, one employs smooth cutoffs by introducing some convergence factors into the integrands of I_1 and I_2 in eq. (5) and extending the integration to infinity. The reason for the violation of condition (6) can be readily understood: The 3-dimensional momentum cutoffs explicitly violate relativistic invariance. Thus, we have encountered a well known problem when the regularization used to make sense out of infinite expressions does not respect a symmetry of the problem. In this situation one may adopt

one of the two different approaches: (1) Continue using the noninvariant regularization but introduce (noninvariant) counter terms which would compensate for this and restore the invariance; (2) Use an invariant regularization. The first approach turns out to be not very useful for the problem under consideration. The reason is that the requirement of the relativistic invariance does not fix the counter terms uniquely as one is always free to add an arbitrary constant to the Lagrangian; this would modify $T_{\mu\nu}^{vac}$ by a term of the form (2). We therefore concentrate on the second approach.

Dimensional regularization. In a d -dimensional spacetime with one time and $d - 1$ space dimensions one has to replace eqs. (1) and (4) by

$$\rho = \frac{1}{2} \mu^{4-d} \int \frac{d^{d-1}k}{(2\pi)^{d-1}} \sqrt{\mathbf{k}^2 + m^2} = \mu^4 \frac{\Gamma(-d/2)}{2(4\pi)^{(d-1)/2} \Gamma(-1/2)} \left(\frac{m^2}{\mu^2} \right)^{d/2}, \quad (10)$$

$$p = \frac{1}{2(d-1)} \mu^{4-d} \int \frac{d^{d-1}k}{(2\pi)^{d-1}} \frac{\mathbf{k}^2}{\sqrt{\mathbf{k}^2 + m^2}} = \mu^4 \frac{\Gamma(-d/2)}{4(4\pi)^{(d-1)/2} \Gamma(1/2)} \left(\frac{m^2}{\mu^2} \right)^{d/2}, \quad (11)$$

where μ is a parameter with dimensions of mass introduced so as to keep the correct dimensions of ρ and p . From the well-known property of the Γ function $\Gamma(-1/2) = -2\Gamma(1/2)$ it immediately follows that eqs. (10) and (11) satisfy the relativistic invariance condition (3). This had to be expected since dimensional regularization respects relativistic invariance. Setting $d = 4 - \epsilon$ one obtains in the limit $\epsilon \rightarrow 0$

$$\rho = -p = -\frac{m^4}{64\pi^2} \left(\frac{2}{\epsilon} + \frac{3}{2} - \gamma - \ln(m^2/4\pi\mu^2) \right) \equiv -\frac{m^4}{64\pi^2} \left(\ln(\Lambda^2/m^2) + \frac{3}{2} \right), \quad (12)$$

where $\gamma \simeq 0.5772$ is the Euler-Mascheroni constant, and in the last equality we traded the parameter μ for a mass scale Λ using the $\overline{\text{MS}}$ renormalization scheme convention. Direct inspection shows that the coefficients of the logarithmically divergent terms in (12) coincide with those obtained using the 3-momentum cutoff.

It should be noted that dimensional regularization is not suitable for studying the power-law divergences since the integrals of the type $\int d^d k (k^2)^\alpha$ vanish identically in this regularization scheme. What we actually learned from the above calculation is that the logarithmically divergent terms in ρ and p indeed satisfy the relativistic invariance condition (3) and that the naive 3-momentum cutoff calculation produced them with correct coefficients.

Four-momentum cutoff. In order to be able to use a relativistically invariant 4-momentum cutoff, one needs

manifestly covariant expressions for ρ and p . One possibility would be to try to convert the integrals over 3-momentum in eqs. (1) and (4) into 4-momentum integrals. However, this procedure is not well defined because the integrals are divergent. To see what kind of difficulties can arise along this route, let us assume that the integrals in (1) and (4) are regulated, and that the regularization (i) either does not modify the integrands or introduces into them an m -independent convergence factor, and (ii) does not affect the covariance properties of the integrals. Under these conditions, one can differentiate the integrals with respect to m^2 . From (1) one then finds

$$\frac{\partial \rho}{\partial m^2} = \frac{1}{2} \int \frac{d^3 k}{(2\pi)^3 2\omega_k}. \quad (13)$$

The integral on the r.h.s. is well known to be relativistically invariant; it can be written in the manifestly invariant form as

$$\frac{\partial \rho}{\partial m^2} = \frac{i}{2} \int \frac{d^4 k}{(2\pi)^4} \frac{1}{k^2 - m^2 + i\varepsilon} = \frac{1}{2} D_F(x, x), \quad (14)$$

where $D_F(x, y)$ is the coordinate-space Feynman propagator of the field ϕ . Integrating eq. (14) one finds, up to an m -independent constant,

$$\rho = \frac{1}{2} \int \frac{d^4 k_E}{(2\pi)^4} \ln \left(1 + \frac{m^2}{k_E^2} \right), \quad (15)$$

where the integral was Wick-rotated to Euclidean space, so that the 4-momentum cutoff can now be utilized. With this cutoff in place, one finds from (15)

$$\rho = \frac{1}{64\pi^2} \left[\Lambda^4 \ln \left(\frac{\Lambda^2 + m^2}{\Lambda^2} \right) + \Lambda^2 m^2 - m^4 \ln \left(\frac{\Lambda^2 + m^2}{m^2} \right) \right] \quad (16)$$

The obtained result appears to be quite sensible: it is just the 1-loop contribution V_1 to the effective potential $V(\phi)$ of a free scalar field. The mass term in the Lagrangian of ϕ can be considered as a tree-level potential $U(\phi) = m^2 \phi^2/2$. The vacuum energy is given by the minimum of the effective potential $V(\phi)$; since for a free scalar field the 1-loop contribution V_1 is ϕ -independent and there are no higher-loop contributions, the vacuum energy just coincides with V_1 .

We shall now show, however, that the differentiation with respect to m^2 that has been used in the derivation of (15) is in general ambiguous and may lead to incorrect results. Comparing the integrands of the right-hand sides of eqs. (1) and (4) and assuming that the differentiation with respect to m^2 is justified, one finds

$$\left(1 - 2m^2 \frac{\partial}{\partial m^2} \right) \rho = (d-1)p. \quad (17)$$

Here we have introduced the factor $d-1$ on the r.h.s. so that the relation is suitable also for the dimensionally-continued case. It is easy to see that in the case of the dimensional regularization the right-hand sides of eqs. (10)

and (11) indeed satisfy (17), and the relativistic invariance condition (3) is also fulfilled. At the same time, if one uses the Euclidean cutoff as a regulator (in conjunction with the differentiation with respect to m^2), eq. (3) is not satisfied. This can be most directly seen by substituting the explicit formula (16) for ρ into (17) and computing the pressure p . Another way to see this is to assume that eq. (3) is satisfied; from (17) one then obtains

$$m^2 \frac{\partial \rho}{\partial m^2} = \frac{d}{2} \rho, \quad (18)$$

which in $d=4$ dimensions has the unique solution $\rho = \text{const} \cdot m^4$, in clear conflict with eq. (16). Thus, eq. (16) cannot be trusted.

The failure of the approach based on the differentiation with respect to m^2 is a consequence of the fact that momentum cutoffs (either 4-dimensional or 3-dimensional) violate one of the two conditions (i) and (ii) discussed above, so that this differentiation is in fact illegitimate. On the other hand, in the case of dimensional regularization both conditions are satisfied; eq. (18) yields $\rho = c_1 \mu^4 (m^2/\mu^2)^{d/2}$ with μ an arbitrary mass parameter and c_1 a (d -dependent) dimensionless constant. This is in accord with eq. (10). Applying dimensional regularization directly to (15) one recovers eq. (10).

Instead of using the differentiation with respect to m^2 in order to arrive at relativistically invariant expressions for the vacuum energy and pressure, one could have started from the effective potential as derived from the functional integral formalism. Since this formalism is manifestly covariant, condition (3) will be automatically satisfied. However, the functional integral formalism yields the effective potential only up to a constant (field-independent) term. While this is quite sufficient when one is looking for the vacuum configuration of the fields, it is not suitable for our purposes since this field-independent constant is exactly what we are seeking. We therefore adopt a different approach.

Let us calculate the trace of the energy momentum tensor of the field ϕ . Using the equation of motion, one formally finds

$$T_\mu^\mu = m^2 \phi^2. \quad (19)$$

In the case of interacting fields this result is modified by quantum anomalies [4], but for the free-field case under consideration it is not anomalous in flat spacetimes. In curved spacetimes, the trace of the energy-momentum tensor is modified by the conformal anomaly [5]. This adds finite terms to the r.h.s. of (19) and thus does not affect the character of the divergences of the vacuum energy; we therefore ignore these terms.

From eq. (19), the vacuum expectation value of T_μ^μ is $\langle 0|T_\mu^\mu|0\rangle = m^2 \langle 0|\phi^2|0\rangle$. On the other hand, from (3) one finds $\langle 0|T_\mu^\mu|0\rangle = \rho - 3p = 4\rho$. This gives

$$\rho = \frac{1}{4} m^2 \langle 0|\phi^2|0\rangle. \quad (20)$$

The vacuum expectation value of ϕ^2 has a well known integral representation, and one finally finds

$$\rho = \frac{m^2}{4} \int \frac{d^4 k}{(2\pi)^4} \frac{i}{k^2 - m^2 + i\epsilon} = \frac{m^2}{4} D_F(x, x). \quad (21)$$

The integral here coincides with the one in eq. (14); however, in the derivation of eq. (21) no differentiation with respect to m^2 has been used. By direct power counting, the leading divergence of the integral in eq. (21) is quadratic. Performing Wick rotation and imposing the Euclidean 4-momentum cutoff one finds

$$\rho = \frac{1}{64\pi^2} \left[\Lambda^2 m^2 - m^4 \ln \left(\frac{\Lambda^2 + m^2}{m^2} \right) \right]. \quad (22)$$

A similar result is obtained if one uses the Pauli-Villars regularization rather than the 4-momentum cutoff [6]. An important consequence of the absence of the quartic divergence in the zero-point energy is that it vanishes for massless fields.

Several comments are in order.

(1) The unrenormalized vacuum energy density ρ that we considered is obviously regularization scheme dependent; therefore the right question to ask is not if one or another scheme results in a quartic divergence, but rather if there exists *any* relativistically invariant regularization that leads to $\mathcal{O}(\Lambda^4)$ terms in ρ . The fact that several regularization schemes leading to quartic divergences in eqs. (1) and (4) violate the relativistic invariance condition (3) does not answer this question because it does not rule out the existence of a such a consistent, relativistically invariant regularization. However, from the representation (20) for the zero-point energy it directly follows by a simple power counting that ρ cannot have quartic divergences.

(2) In our discussion we considered flat spacetimes. In a curved spacetime, the curvature only influences the modes with the wavelength $k^{-1} \gtrsim a$ where a is the local curvature radius, as one can always find a local Lorentz frame in which the spacetime is flat in the region of the size $r \lesssim a$. Since the ultraviolet divergences correspond to the smallest wavelengths with $k^{-1} \ll a$, their nature does not depend on whether the spacetime is flat or curved. Therefore the divergence of the vacuum energy is unaffected by the spacetime curvature.

(3) In models with broken supersymmetry the vacuum energy is known to have no quartic divergence as a result of the cancellations between the contributions of the fermionic and bosonic superpartners. We have shown here that in fact the quartic divergences are absent even without supersymmetry since the relativistic invariance precludes their appearance in the *individual* contributions of different fields.

(4) We have considered the zero-point energies of free fields. As was pointed out above, in the interacting case the trace formula (19) is modified by the flat-space trace anomaly [4]; therefore this case requires a special study.

(5) Although our results have not led to a solution of the cosmological constant problem, we believe that they can be relevant to it. From the absence of the quartic divergences it follows that in flat spacetimes the zero-point energies of free massless fields vanish. In particular, the zero-point energy of the free electromagnetic field is zero [7]. In a curved spacetime of curvature radius a the vacuum energy of a free massless field is of order a^{-4} ; for the present-day universe this is vanishingly small compared to the critical energy density ρ_c . Thus, free massless fields give negligible contributions to the vacuum energy.

Acknowledgements. The author is grateful to M.C. Bento, Z. Berezhiani, H.B. Nielsen, L.B. Okun, M.E. Shaposhnikov and A.Yu. Smirnov for useful discussions, and to V.A. Rubakov and J. Smit for useful correspondence. He was supported by the Calouste Gulbenkian Foundation as a Gulbenkian Visiting Professor at Instituto Superior Técnico.

-
- [1] S. Perlmutter *et al.* *Astrophys. J.* **517**, 565 (1999); A. G. Riess *et al.* *Astron. J.* **116**, 1009 (1998).
 - [2] For reviews see, e.g., S. Weinberg, *Rev. Mod. Phys.* **61**, 1 (1989); V. Sahni and A. Starobinsky, *Int. J. Mod. Phys. D* **9**, 373 (2000); S. M. Carroll, *Living Rev. Rel.* **4**, 1 (2001).
 - [3] There are several ways to arrive at this result. One may note that the vacuum can be considered a perfect fluid; using the general form of the energy-momentum tensor of perfect fluids $T_{\mu\nu} = -pg_{\mu\nu} + (\rho + p)U_\mu U_\nu$ where U_μ is the four-velocity vector, one obtains (3). Alternatively, one may start from a flat spacetime with Minkowski metric $g_{\mu\nu} = \eta_{\mu\nu} \equiv \text{diag}(1, -1, -1, -1)$ so that eq. (3) is automatically satisfied, and then notice that any curved spacetime is locally flat.
 - [4] C. G. Callan, S. R. Coleman and R. Jackiw, *Annals Phys.* **59**, 42 (1970).
 - [5] D. M. Capper and M. J. Duff, *Nuovo Cim. A* **23**, 173 (1974); S. Deser, M. J. Duff and C. J. Isham, *Nucl. Phys. B* **111**, 45 (1976).
 - [6] Note that the Pauli-Villars regularization cannot be applied directly to eq. (1) because when the regularization is lifted (i.e. in the limit of the infinite masses of the regulator fields) the original divergent expression is not recovered in that case.
 - [7] This does not apply to the zero-point energies inside cavities or when the boundaries are present, and so the Casimir energies do not vanish, of course. However, when calculated in a covariant way and with a relativistically invariant regularization, these energies must be finite, and no subtraction of the energy of the free vacuum is necessary.